



Increased Expression of the lnc H19 Gene in the Plasma of People with Breast Cancer

Kazhaleh Mohammadi^{1*}, Sadegh Shojaei Baghini², Mohammad Ali Saremi³

¹ Researcher and lecture assistant at the Medical Laboratory Science Department, Knowledge University, Kurdistan Region, Erbil, Iraq.

² Plant Biotechnology, Agriculture Biotechnology Department, National Institute of Genetics Engineering and Biotechnology, Tehran, Iran.

³ Personalized Medicine Research Center of AmicitGen, Tehran, Iran.

DOI: 10.22034/PMJ.2020.43454

*Corresponding author: kazhaleh Mohammadi, Researcher and lecture assistant at the Medical Laboratory Science Department, Knowledge University, Kurdistan Region, Erbil, Iraq. Email: Kazhaleh.mohammadi@Knu.edu.iq

Submitted: 2020-04-25

Accepted: 2020-05-17

Keywords:

lncRNAs
Breast cancer
Gene expression
Plasma
qRT-Real-Time PCR

©2020. Personalized Medicine Journal

Abstract:

The majority of ncRNAs are known as long non-coding RNAs (lncRNAs) whose length exceeds 200 nucleotides. H19, a lncRNA, is the transcription product of the H19 gene, an oncogene in breast cancer, and is highly expressed in cancer tissues compared with normal tissues. The expression level of H19 is associated with the oncogenesis, proliferation, invasion, metastasis, and drug resistance of breast cancer. H19 expression levels were detected in breast cancer plasma using qRT-Real-Time PCR assay in 50 breast cancer samples and 50 healthy control samples. The results showed that the expression of this gene in both the tissue and the plasma of patients increased compared to that of healthy individuals.

INTRODUCTION

The lnc H19 gene is located in an imprinted region of chromosome 11 near the insulin-like growth factor 2 (IGF2) gene. This gene is only expressed from the maternally-inherited chromosome (1). The product of this gene is a long non-coding RNA which functions as a tumor suppressor. H19 promotes biological processes such as apoptosis, angiogenesis, inflammation, and cell death. Furthermore, gene ontology (GO) analyses predict that H19 is connected with neurogenesis, angiogenesis, and inflammation through DNA transcription, RNA folding, methylation, and gene imprinting (2). The aberrant expression of H19 is associated with multiple diseases, including carcinoma, sarcoma, type 2 diabetes, and hypertrophic cardiomyopathy (3). Abnormal expression of H19 has been demonstrated in a variety of cancer cells, such as gastric cancer, pancreatic cancer, liver cancer and breast cancer cells, affecting the development and progression of cancer through various mechanisms (4). H19 is an estrogen-regulated transcript, and a wide array of studies have confirmed that it regulates cell differentiation and proliferation. In addition, research into its application in the clinical setting has revealed that H19 has the potential to serve as a biomarker for cancer and shows promise as a novel therapeutic option in oncology (5). Breast cancer is a malignant tumor deriving from the epithelium of the mammary gland. Metastasis is the leading cause of mortality in patients with breast cancer. The H19 gene is an oncogene in breast cancer and is highly expressed in cancer tissues compared with normal tissues (6). The expression level of H19 is associated with the oncogenesis, proliferation,

invasion, metastasis, and drug resistance of breast cancer, but the underlying molecular mechanisms differ greatly (7). The molecular mechanisms underlying H19-associated carcinogenesis may involve several aspects. The H19 promoter is activated by the E2F transcription factor 1 (E2F1), which promote cell cycle progression (particularly in the S-phase) of MCF-7 cells. H19 knockdown increases the DNMT3B-mediated methylation of Nctc1, a gene encoding lncRNAs, within the Igf2-H19-Nctc1 locus (8). Thus, H19 alters DNA methylation and leads to breast tumorigenesis. A new lncRNA within the H19/IGF2 locus known as 91H is an antisense gene to H19. Furthermore, it has been shown to regulate H19 and IGF2 expression levels by epigenetic modifications and increase the tumorigenic properties of MDA-MB-231 cells both in vitro and in vivo (9). In summary, H19 plays an oncogenic role in breast tumorigenesis. Serum H19 levels may possess clinical significance in the early diagnosis, treatment, and prognosis of breast cancer. The current study investigated whether H19 is detectable in the plasma of breast cancer patients compared with matched healthy controls using qRT-Real-Time PCR assay.

METHODS AND MATERIALS

Our subjects consisted of 40 individuals with breast cancer that their breast tissue was stored in Breast Cancer Tissue Bank of Emam Khomeini hospital Tehran, Iran between November 2012 and October 2013. All this patients were clinically diagnosed according to the typical signs of breast cancer criteria, which were obtained from histopathological examination by expert

oncologists. All patients in this study were not subjected to any preoperative radiotherapy or chemotherapy; they had no other malignant diseases or injuries before sample collection. The informed consent was obtained from all patients who were enrolled in this study. Also, we selected 50 controls that matched to patients with respect to age, place of residence and race/ethnicity. The informed consent was obtained from all patients and healthy who were enrolled in this study. The patients' ages ranged from 47 to 72 years, and the median was 59 years. The healthy participants had not been exposed to any potentially harmful chemicals and had no malignant diseases, acute diseases, or injuries at the time of sample collection. The ages of healthy participants ranged from 29 to 66 years with the median age of 47 years. RNA samples were obtained from samples using by RNeasy FFPE Kit according to manufacturer's instruction (Cat. No. / ID: 73504). Reverse transcription reaction was carried out using PrimeScript™ RT reagent kit with gDNA eraser (Takara, Japan) in accordance with the manufacturer's instructions. Real-time PCR was then performed using RealQ Plus Master mix green (Ampliqon, Denmark) as implemented in Rotorgene Corbett 6000 Real-Time. The sequences of the PCR primers for H19 and β -actin are shown in Table 1. β -actin was selected as an internal control.

The expression level of H19 was obtained using the $2^{-\Delta\Delta Ct}$ method.

Table 1. Primer Sequences

Primer	Sequence
Lnc H19	F: 5'-TGCTGCACTTTACAACCACTG-3' R: 5'-ATGGTGCTTTGATGTTGGGC-3'
B-actin	F: 5'-TCCTCTCCCAAGTCCACACA-3' R: 5'-GCACGAAGGCTCATCATTCA-3'




Fig 1. β -actin was selected as an internal control

RESULTS

In this study, qRT-Real-Time PCR was performed to analyze 40 breast cancer and 50 control plasma samples. It was found that H19 expression was significantly higher in breast cancer plasma relative to the control plasma (Fig. 1, $p = 0.004$). The data indicates that H19 is overexpressed in the circulating plasma of human breast cancer patients.

DISCUSSION

The majority of ncRNAs are known as long non-coding RNAs (lncRNAs) whose length exceeds 200 nucleotides. LncRNAs were first regarded as the "noise" of gene transcription (10). Because of their position relative to the protein-coding genes, lncRNAs can be roughly divided into antisense lncRNAs, enhancer lncRNAs, large intergenic non-coding RNAs, bidirectional lncRNAs, and intronic transcript (11). LncRNAs can regulate gene expression via multiple mechanisms at the epigenetic, transcriptional, and post-transcriptional levels. They participate in the regulation of a variety of cell activities, such as cell differentiation, proliferation, invasion, apoptosis, and autophagy (12) by interacting with RNAs, DNAs, or proteins. The H19 gene is located in 11p15.5, an imprinted region of chromosome 11, adjacent to the insulin-like growth factor 2 (IGF2) gene and is expressed only from the maternally inherited chromosome; IGF2, however, is expressed only from the paternally inherited chromosome (13). H19, a lncRNA, is the transcription

product of the H19 gene, and diversified transcript variants exist due to alternative splicing (14). A recent case-control study in China revealed that high H19 expression levels were associated with an increased risk of breast carcinogenesis in both codominant and dominant models, and the association was more apparent in patients with estrogen receptor-positive (ER+), human epidermal growth factor receptor 2-negative (HER2-), and ER+-HER2-negative (HER2-) molecular subtypes (15). Numerous studies have indicated that circulating nucleic acids are detectable in the human peripheral circulation system. Matouk et al. indicated that H19 is highly expressed in all common metastatic sites, regardless of primary tumor origin, and its expression is tightly correlated with BC metastatic potential (1-17). Vennin et al. demonstrated that the overexpression of H19/miR-675 plays a pivotal role in enhancing the proliferation and migration of BC cells and increasing tumor growth and metastasis (18). Therefore, the detection of circulating nucleic acids in plasma or serum could be utilized as a non-invasive diagnostic tool for human cancer (19-20). The current study evaluated changes in lncRNA H19 gene expression in the plasma of breast cancer patients as a non-invasive biomarker. The results showed that H19 expression was significantly higher in breast cancer plasma relative to control plasma ($p = 0.004$). The data indicates that H19 is overexpressed in the circulating plasma of human breast cancer cases. The results of this study are consistent with the results of the study by Zhang et al., who evaluated the expression of the lncH19 gene in the tissue and plasma of breast cancer patients (21). Their results indicated that the expression of this gene in both the tissue and the plasma of patients was increased compared to that in healthy individuals. It is recommended that this study be performed on a larger scale and that the results be compared with the clinical characteristics of individuals.

REFERENCE

- Kapranov P, Cheng J, Dike S, Nix DA, Duttgupta R, Willingham AT, Stadler PF, Hertel J, Hackermuller J, Hofacker IL, et al: RNA maps reveal new RNA classes and a possible function for pervasive transcription. *Science* 316:1484-1488,2007.
- Dahariya S, Paddibhatla I, Kumar S, Raghuvanshi S, Palapati A and Gutti RK: Long non-coding RNA: Classification, biogenesis and functions in blood cells. *Mol Immunol* 112: 82-92, 2019.
- M.J. Duffy, D. Evoy and E.W. McDermott, CA 15-3: uses and limitation as a biomarker for breast cancer, *Clin Chim Acta* 411 (2010), 1869-1874.
- N. Patani, L.A. Martin and M. Dowsett, Biomarkers for the clinical management of breast cancer: international perspective, *Int J Cancer* 133 (2013), 1-13.
- S. van Heesch, M. van Iterson, J. Jacobi, et al., Extensive localization of long noncoding RNAs to the cytosol and monoand polyribosomal complexes, *Genome Biol* 15 (2014), R6.
- Raveh E, Matouk IJ, Gilon M and Hochberg A: The H19 Long non-coding RNA in cancer initiation, progression and metastasis-a proposed unifying theory. *Mol Cancer* 14:184,2015.
- Yoshimura H, Matsuda Y, Yamamoto M, Kamiya S and Ishiwata T: Expression and role of long non-coding RNA H19 in carcinogenesis. *Front Biosci (Landmark Ed)* 23: 614-625, 2018.
- Bao MH, Szeto V, Yang BB, Zhu SZ, Sun HS and Feng ZP: Long non-coding RNAs in ischemic stroke. *Cell Death Dis* 9: 281, 2018.
- Zeng Y, Li TL, Zhang HB, Deng JL, Zhang R, Sun H, Wan ZR, Liu YZ, Zhu YS and Wang G: Polymorphisms in IGF2/H19 gene locus are associated with platinum-based chemotherapeutic response in Chinese patients with epithelial ovarian cancer. *Pharmacogenomics* 20: 179-188, 2019.
- Ghaedi H, Zare A, Omrani MD, Doustimotlagh AH, Meshkani R, Alipoor S and Alipoor B: Genetic variants in long noncoding RNA H19 and MEG3 confer risk of type 2 diabetes in an Iranian population. *Gene* 675: 265-271, 2018.
- N. Giannoukakis, C. Deal, J. Paquette, et al., Parental genomic imprinting of the human IGF2 gene, *Nat Genet* 4 (1993), 98101.
- T. Dugimont, J.J. Cury, N. Wernert, et al., The H19 gene is expressed within both epithelial and stromal components of human invasive adenocarcinomas, *Biol Cell* 85 (1995), 117124.
- E. Adriaenssens, S. Lottin, T. Dugimont, et al., Steroid hormones modulate H19 gene expression in both mammary gland and uterus, *Oncogene* 18 (1999), 4460-4473.
- M. Zhu, Q. Chen, X. Liu, et al., lncRNA H19/miR-675 axis represses prostate cancer metastasis by targeting TGFBI, *FEBS J* 281 (2014), 3766-3775.
- M. Zhu, Q. Chen, X. Liu, et al., lncRNA H19/miR-675 axis represses prostate cancer metastasis by targeting TGFBI, *FEBS J* 281 (2014), 3766-3775.
- M. Zhu, Q. Chen, X. Liu, et al., lncRNA H19/miR-675 axis represses prostate cancer metastasis by targeting TGFBI, *FEBS J* 281 (2014), 3766-3775.
- Y.J. Jiang and D.D. Bikle, LncRNA: a new player in $1\alpha, 25(\text{OH})_2$ vitamin D(3) /VDR protection against skin cancer formation, *Exp Dermatol* 23 (2014), 147-150.
- Z.F. Yang, P. Ngai, D.W. Ho, et al., Identification of local and circulating cancer stem cells in human liver cancer, *Hepatology* 47 (2008), 919-928.
- Zhou W, Ye XL, Xu J, Cao MG, Fang ZY, Li LY, Guan GH, Liu Q, Qian YH and Xie D: The lncRNA H19 mediates breast cancer cell plasticity during EMT and MET plasticity by differentially sponging miR-200b/c and let-7b. *Sci Signal* 10: pii: eaak9557,2017.
- Zhou W, Ye XL, Xu J, Cao MG, Fang ZY, Li LY, Guan GH, Liu Q, Qian YH and Xie D: The lncRNA H19 mediates breast cancer cell plasticity during EMT and MET plasticity by differentially sponging miR-200b/c and let-7b. *Sci Signal* 10: pii: eaak9557, 2017
- De Martino M, Forzati F, Marfella M, Pellicchia S, Arra C, Terracciano L, Fusco A and Esposito F: HMGA1P7-pseudogene regulates H19 and Igf2 expression by a competitive endogenous RNA mechanism. *Sci Rep* 6: 37622, 2016.