



## Kefir's Hidden Arsenal: Examining the Effect of Lactobacilli Supernatant on Antibiotic Resistance Genes and Virulence in *Salmonella Typhimurium*

Salman Imran Hammadi<sup>1</sup>, Zahra shafiei<sup>2\*</sup>, Bizhan Nemanpour<sup>3</sup>

<sup>1</sup>Department of Microbiology, School of Basic Sciences, Science and Research Branch, Islamic Azad University, Tehran, Iran.

<sup>2</sup>Department of Biology, School of Science and Agriculture, Islamic Azad University, Roudehen Branch, Tehran, Iran.

<sup>3</sup>Department of Microbiology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran.

DOI: 10.22034/pmjournal.2023.2018309.1024

\*Corresponding author: Zahra shafiei, Department of Biology, School of Science and Agriculture, Islamic Azad University, Roudehen Branch, Tehran, Email: z.shafiee@riau.ac.ir.

Submitted: 2023-09-19

Accepted: 2023-11-30

### Keywords:

Salmonella

*S. typhimurium*

Lactobacillus

Kefir

Antibiotic resistance

©2023. Personalized Medicine Journal

### Abstract:

Salmonella, a prominent foodborne pathogen, poses significant health risks, causing both intestinal and extra-intestinal infections. Recognizing the potential of lactobacilli as probiotics due to their ability to produce substances inhibiting multidrug-resistant bacteria, this study aimed to assess antibiotic resistance, pathogenic gene frequency, antibacterial effects of lactobacillus supernatant from kefir, and its impact on resistance and pathogenicity gene expression.

In Tehran hospitals, 150 isolates from 240 clinical samples were collected and identified as *Salmonella typhimurium* using biochemical and serotype tests. Antibiotic sensitivity was assessed, and the frequencies of antibiotic resistance genes (*tetA*, *tetB*, and *floR*) and pathogenicity genes (*sipA*, *spvC*, and *invA*) were investigated. Lactobacilli from kefir were isolated, and the minimum inhibitory concentration of lactobacillus supernatant was determined. The relationship between supernatant treatment and *tetA* and *sip* gene expression was examined using Real-time PCR.

Results revealed 38% of strains as *Salmonella typhimurium* serotype, displaying high resistance to ampicillin, tetracycline, and nitrofurantoin. Pathogenicity genes *invA* and *sip* exhibited high frequencies of 100% and 70.2%, respectively. Lactobacillus supernatant showed an MIC of 80 µg/ml, effectively reducing *tetA* and *sip* gene expression by 42.2% and 55.7%, respectively.

In conclusion, the study underscores the high antibiotic resistance in *Salmonella typhimurium* and suggests Meropenem, Trimethoprim Sulfamethoxazole, and Ampicillin-Sulbactam as effective treatments. Moreover, lactobacillus supernatant demonstrated significant potential against *Salmonella typhimurium*, highlighting lactobacilli as promising probiotics. This health-oriented strategy presents a viable solution for treating Salmonella infections and preventing their spread.

## INTRODUCTION

Enterobacteriaceae family is a diverse and extensive group of Gram-negative bacilli encompassing a wide range of species and genera. The most important pathogenic genera in this family that affect humans include *Escherichia* spp, *Klebsiella* spp, *Salmonella* spp, *Shigella* spp, and *Yersinia* spp (1). This large and heterogeneous family produces a set of antigenic structures and virulence factors that cause disease in humans and animals (2). *Salmonella* species are regarded as common foodborne infections due to their vast range of animal reservoirs, causing considerable global public health issues with significant economic

repercussions. These gram-negative bacteria are rod-shaped, small, and lack capsules, but have surrounding flagella that measure 4.5-2.5 microns. Over 2700 unique *Salmonella* species have been found to date in diverse places throughout the world (3, 4).

It is estimated that 16 million cases of typhoid fever, 1.3 million cases of gastroenteritis, and 3 million deaths worldwide are attributed to salmonella. Studies have demonstrated that poultry serves as a significant reservoir of human salmonellosis. This is due to the bacterium's ability to colonize the digestive system and cloaca of poultry, which are important sites for salmonella colonization. (5-7).

*Salmonella enteritidis* and *Salmonella typhimurium* have been reported as two common serotypes isolated from different cases. Infection caused by these two requires treatment or even leads to hospitalization (8). The rapid emergence of antimicrobial resistance by microbial pathogens is a threat to public health. The emergence of microbial resistance is not a new phenomenon and the production of new antibiotics has become a challenge in disease control in poor and developing countries. In 2014, the World Health Organization (WHO) named antibiotic resistance as a major global threat. This organization has reported the increase of drug resistance in all parts of the world by reviewing the statistics of 114 countries (9). *Salmonella* species employ genes and pathogenic determinants for the invasion of host cells and the initiation of pathogenic processes. The *invA* gene is responsible for encoding the invasive protein *invA*, which plays a pivotal role in attacking intestinal epithelial cells, ultimately resulting in the onset of the disease. Additionally, this gene contributes to the transportation of the *invE* protein. The *invA* gene facilitates the ingress of bacteria into epithelial cells and exhibits structural similarities to genes associated with flagella biosynthesis. Furthermore, the *SipA* protein assumes a crucial function in actin polymerization. *Salmonella* employs the type 3 secretion system (T3SS) to inject the 12 effector proteins encoded by the pathogenicity island 1 (SPI-1) into the host cell. These proteins, likely the first to be released by *SipC* and *SipA*, include *SipA* and *SipC*. The *SipA* and *SipC* proteins induce membrane roughness, promote invasion, and contribute to stabilization through their direct interaction with the actin cytoskeleton. Additionally, they play a regulatory role in actin movement and dynamics (10-12).

According to different research and publications, antibiotic resistance in *Salmonella* is becoming an increasing concern. Antibiotic-resistant nontyphoidal *Salmonella* infections are on the rise, according to the Centers for Disease Control and Prevention (CDC), with some strains resistant to key medicines such as ciprofloxacin, azithromycin, and ceftriaxone. *Salmonella* strains have also been reported to

be resistant to a variety of antibiotics, including streptomycin, gentamicin, and sulfadimethoxine. Furthermore, a strain of extremely drug-resistant (XDR) *Salmonella typhi* has evolved that is resistant to all antibiotic classes except two (13). This trend emphasizes the necessity of combating antibiotic resistance in *Salmonella* with comprehensive tactics, such as the investigation of alternate control measures and the identification of efflux pumps, regulators, and inhibitors to tackle multidrug resistance.

Probiotics are live microorganisms that are similar to the beneficial microorganisms found in the human gut (14). Probiotics have been demonstrated to effectively alleviate symptoms of lactose intolerance by breaking down lactose with the production of the beta-D-glucosidase enzyme, thereby, preventing or minimizing occurrences of diarrheal diseases. Additionally, probiotics play a role in preventing and managing allergies, as evidenced by studies indicating that probiotics containing *Lactobacillus GG* may reduce the prevalence of atopic eczema in later stages of life. Beyond these applications, probiotics exhibit important properties such as anti-genotoxic, anti-mutagenic, and anti-cancer effects, along with a reduction in the production of carcinogenic or toxic metabolites. Epidemiological studies further support the use of probiotics in reducing the incidence of colon cancer through various mechanisms (15-17).

Kefir is a fermented milk beverage made by lactic acid bacteria, acetic acid bacteria, and yeast. *Lactobacilli* are among the most often encountered bacteria in kefir (18, 19). These microorganisms have been shown to reduce bacteria pathogenicity and hinder several virulence factors (20). The aim of this study is to investigate the impact of *Lactobacilli* Supernatant on antibiotic resistance genes and virulence in *Salmonella Typhimurium*.

## MATERIALS AND METHODS

This research was descriptive-cross-sectional. 240 stool samples from patients with diarrhea were collected from Imam Khomeini, Shahada Tajrish and Luqman Hospitals during the period of 1401 to 1402

**Table 1.** The function of virulence genes in the pathogenic process of *Salmonella* bacteria

Gene	Gene function and role
InvA	Role in the invasion and invasion of intestinal mucosa and epithelial tissue
SipA	Actin polymerization, role in bacterial motility and spread
SpvC	<ol style="list-style-type: none"> <li>Helping the survival of bacteria inside the host cells</li> <li>Helping the growth and reproduction of the pathogen in the host's body and in extra-intestinal places</li> <li>Helping the systemic spread of the pathogen in the host's body (systemic infection)</li> </ol>

in Tehran and were transferred to the laboratory at 4°C using the appropriate method. The strains were cultured in TSB medium and then it was transferred to EMB and McConkey agar medium, warm staining was done from suspicious colonies and finally they were subjected to biochemical tests such as Oxidase, ONPG, MR-VP for final confirmation (21). For drug sensitivity determination, Mueller Hinton Agar culture medium used in performing the disk diffusion test by McFarland half solution for 20-24 hours (22). Based on the Kaufman-White method, the serotypes of Salmonella isolates can be identified, this method is based on the agglutination of bacteria with specific somatic (O) and flagella (H) antisera (23). After drug resistance, in order to check the frequency of antibiotic resistance genes and pathogenicity genes, molecular tests were performed. The existence of these genes was investigated by multiplex PCR against the antibiotic resistance genes *tetA*, *tetG*, and *floR* and the pathogenicity genes *sipA*, *spvC*, and *invA* and Gel Electrophoresis was done (24) (Table 2).

The bacteria isolated from kefir were cultured and their extracts were obtained. In this research, microscopic and physicochemical methods were used for the initial identification of the obtained strain (25). The effects of these extracts on the selected strain were performed using the microdilution method, and then the effect of this treatment on the expression of the pathogenic gene *sip* and the resistance gene to the antibiotic tetracycline (test) was done using Real-time PCR method. The results are interpreted using the  $\Delta$ CT

**Table 2.** Sequences of primers used in this research

desired gene	Primer sequence	PCR product size
floR	AACCCGCCCTCTGGATCAAGTCAA CAAATCACGGGCCACGCTGTATC	548
tetA	GCT ACA TCC TGC TTG CCT TC CAT AGA TCG CCG TGA AGA GG	210
tetB	TTG GTT AGG GGC AAG TTT TG GTA ATG GGC CAA TAA CAC CG	659
invA	CGCGCCCCGATTTTCTCTGGA AATGCGGGGATCTGGGCGACAAG	321
sipB/C	ACAGCAAATGCGGATGCTT GCGCGCTCAGTGTAGGACTC	232
spvC	ACTCCTTGCAACAACCAAATGCGGA TGTCTTCTGCATTTCCACCACATCA	424

**Table 3.** Information of the strains isolated

Total number of 150 pieces	Number of strains		Serotypes
	Female (total number 58)	Male (total number 92)	
57 (38 percent)	21	36	<i>Salmonella typhimurium</i>
93 (62 percent)	37	56	<i>Salmonella enteritidis</i>


method. The quantification of the augmentation in gene copy numbers of specific genes in the presence of nanoparticles was computed by dividing the number of gene copies post-treatment by the number of copies pre-treatment. The assessment of resistance is contingent upon this observed augmentation, a parameter that varies according to the specific antimicrobial substance under consideration.

All the methods were presented in supplementary file.


## RESULTS

In this research, 150 isolates isolated from people suspected of intestinal disease from Imam Khomeini, Shahada Tajrish, and Luqman hospitals in Tehran were prepared and then subjected to morphological, biochemical, and antibiotic resistance studies by disc diffusion method, and of these, 57 samples The strain that was known as Salmonella typhimurium. (Table3). The frequency of resistance gene to tetracycline (*tetA/B*) and florfenicol (*floR*) antibiotics were checked by PCR method. Then the lactobacillus isolated from the cultured kefir and its supernatant were obtained and treated against the selected strain, and finally, changes in the expression of the pathogenicity gene (*sipA*) and the tetracycline antibiotic resistance gene (*tetA*) were determined using the Realtime PCR method (Figure 1-2).

The Salmonella strains in this study showed high antibiotic resistance, so that compared to ampicillin (100%), tetracycline (100%) and nitrofurantion (84.2



**Fig1.** Electrophoresis results of PCR products for *tetA*, *tetB* and *floR* genes, for isolates 1 to 57, M gene marker, + positive control (*Salmonella typhimurium* ATCC 14028 strain) and - negative control (distilled water).



**Fig 2.** Electrophoresis results of PCR products for *sipA*, *invA* and *spvC* genes, for isolates 1 to 57, M gene marker, + positive control (*Salmonella typhimurium* ATCC 14028 strain) and - negative control (distilled water).

to 100%) the most showed resistance and compared to meropenem (1.1 to 1.8 percent), trimethoprim sulfamethoxazole (1.8 to 2.2 percent) and ampicillin-

sulbactam (3.5 to 4.3 percent) Showed. As can be seen, *Salmonella Typhimurium* serovar also showed high resistance to different classes of antibiotics,

showing the highest resistance to ampicillin (100%), tetracycline (100%) and nitrofurantion (84.2%). They also showed sensitivity to the antibiotic meropenem (1.8%), trimethoprim-sulfamethoxazol (1.8%), and ampicillin-sulbactam (3.5%)(Table 4).

The genetic data results indicate that the tetracycline resistance gene was most prevalent, with 86% of the strains carrying the gene. Additionally, a high frequency of the florfenicol resistance gene was observed, with 56.1% (32) of the strains containing this None of the strains contained all three studied genes, but 54.4% (31) of the strains contained both tetracycline and florfenicol resistance genes (Table 5).

The results of the genetic data show that the highest frequency of the pathogenicity gene in the studied

strains was observed for the *invA* gene and all the strains (100%) had this gene, the frequency of the sip gene was 72.2% (40) and None of the strains (0%) had the *spvC* gene (Table 6).

The results showed that bacterial growth was inhibited at a concentration of 160 µg/ml of the supernatant, and this concentration was considered as MIC, and a concentration of 80 µg/l was reported as sub-MIC (Table 7).

Results of tetA gene expression change in pre and post treatment with Lactobacillus supernatant by Real time PCR method evaluated by melting curves and  $\Delta\Delta CT$  Method. The expression level, which is a measure of  $\Delta\Delta CT$ , is obtained using the calculations of cycle thresholds (Ct) finally, to calculate the fold change,

**Table 4.** The results of the frequency of antibiotic resistance of isolates

SALMONELLA ENTERITIDIS (93)			SALMONELLA TYPHIMURIUM(57)		
Antibiotics	Percent	resistant	Percent	Number	DENSITY
Ampicillin	% 100	93	% 100	57	10
Ampicillin-sulbactam	% 4/3	4	% 3/5	2	100-10
Amoxicillin-clavulanic acid	% 69/9	65	% 54/4	31	20-10
Cefataxime	% 51/8	54	% 31/6	18	30
Amy Panam	% 25/8	24	% 36/8	21	10
Gentamicin	% 6/5	6	% 5/3	3	10
Tetracycline	% 100	93	% 100	57	30
Ciprofloxacin	% 22/6	21	% 14	8	5
Nalidixic acid	% 38/7	36	% 43/9	25	30
Trimethoprim Sulfomethoxazole	% 2/2	2	% 1/8	1	25
Chloramphenicol	% 22/6	21	% 7	4	30
Azithromycin	% 25/8	24	% 26/3	15	300
Nitrofurantion	% 100	193	% 84/2	48	300
Meropenem	% 1/1	1	% 1/8	1	2

**Table 5.** Frequency of antibiotic resistance genes in Salmonella typhimurium strains

PERCENT	NUMBER	GENE NAME
% 86	49	TETA
% 1/8	1	TETB
% 56/1	32	FLOR
% 0	0	ALL GENES
% 54/4	31	TET+FLOR

**Table 6.** Frequency of sip, invA and spvC pathogenicity genes in Salmonella Typhimurium strains

PERCENT	NUMBER	GENE NAME
% 0	0	SPVC
% 100	57	INVA
% 70/2	40	SIP

**Table 7.** The results of the minimum inhibitory concentration of treatment with lactobacillus supernatant against the selected strain

CONCENTRATION OF TUNGSTEN OXIDE NANOPARTICLES (MICROGRAMS/ML)	SALMONLATIFI MORIUM
160	MIC
80	SUB MIC
2560	INITIAL CONCENTRATION

the expression of the treated gene (0.579012607) should be subtracted from the control (1.00126) ) to be calculated, which was calculated as 0.578 in this research, which shows that the expression of resistance genes decreased by 42.2% during the treatment. As shown in Table 6-4, the statistical analysis of the obtained results was done and the relationship between the treatment and the obtained results was tested under the condition of  $P < 0.05$  and the results were confirmed. (Table 8) In order to calculate the average relative expression, the average of  $\Delta\Delta CTs$  was first taken and the result was calculated with the formula of relative expression (Table 9) (Figure 3).

Results of sip gene expression change in pre and post treatment with Lactobacillus supernatant by Realtime PCR method evaluated by melting curves and  $\Delta\Delta CT$  Method, the expression level, which is a measure of  $\Delta\Delta CT$ , to calculate the fold change, the expression of the treated gene (0.443943) should be subtracted from the control (1.00098). in this study it was calculated as 0.443508, which shows that the expression of

resistance genes decreased by 55.7% during the treatment. the statistical analysis of the obtained results was done and the relationship between the treatment and the obtained results was tested under the condition of  $P < 0.05$  and the results were confirmed. In order to calculate the average relative expression, the average of  $\Delta\Delta CTs$  was first taken and the result was calculated with the formula of relative expression (Table 10, 11) (Figure 3).

## DISCUSSION

The research findings indicate that Salmonella typhimurium constitutes a significant portion (38%) of isolates in individuals with intestinal infections. Notably, 63% of these isolates are found in men, emphasizing the prevalence of this strain in the male population. Antibiotic resistance patterns reveal high resistance to ampicillin, tetracycline, and nitrofurantion, with multidrug resistance observed. Meropenem, trimethoprim-centosulfamethoxarol, and amoxicillin-sulbactam are suggested for treatment.

**Table 8.** Mean relative expression for tetA gene

STANDARD DEVIATION (SD)	MEAN RELATIVE EXPRESSION
0/061826	1 Control
0/01853	0/579012607 treatment

**Table 9.** Mean relative expression for sip gene

STANDARD DEVIATION (SD)	MEAN RELATIVE EXPRESSION
0/0537	1 Control
0/04601	0/44237 treatment




Fig 3. Mean relative expression plot for tetA and sip gene

Table 10. Calculation of relative expression and statistical analysis of data obtained from Real-time PCR reaction for tetA gene

House (16S rRNA)			
Name	CT	CT	Average ( CT )
Treat	25.71	25.53	25.62
Non-Treat	25.91	25.88	25.895

Gene (tetA)			
Name	CT	CT	Average ( CT )
Sample 1 Treat	26.29	26.23	26.26
Sample 2 Treat	26.41	26.37	26.39
Sample 3 Treat	26.16	26.09	26.125
Sample 1 Non-Treat	25.89	25.63	25.76
Sample 2 Non-Treat	25.87	25.78	25.825
Sample 3 Non-Treat	25.61	25.69	25.65

ΔΔCT Method			
Name	ΔCT	ΔΔCT	Fold change
Sample 1 Treat	0.64	0.775	0.584388624
Sample 2 Treat	0.77	0.84	0.558643569
Sample 3 Treat	0.505	0.75	0.594603558
Sample 1 Non-Treat	-0.135	0.015	
Sample 2 Non-Treat	-0.07	0.08	
Sample 3 Non-Treat	-0.245	-0.095	
<b>Avg Fold change</b>	<b>0.5792119</b>		

T-Test	
P-value	Significant or Not

Genetic analysis indicates a high frequency of tetracycline resistance genes and pathogenicity genes, particularly *invA*. The inhibitory potential of lactobacillus supernatant against Salmonella is demonstrated, leading to decreased expression of tetracycline resistance and sip pathogenicity genes.

Comparing with other studies, Nosrati et al. (2012) found a high prevalence of Salmonella typhimurium and Salmonella enteritidis in food samples, emphasizing the risk from food sources in salmonella infections (26). Nazari Moghadam et al. (2023) aimed to investigate the prevalence of virulent antibiotic-resistant Salmonella spp. strains in Iranian poultry

markets, highlighting the importance of monitoring this pathogen in food sources (27). Ribeiro et al. (2011) observed an increase in antimicrobial resistance among Salmonella Enterica, emphasizing the impact on public health (28). Dantas et al. (2020) highlighted the persistence and pathogenic potential of Salmonella in poultry slaughterhouse, indicating resistance to tetracycline and the role of biofilm production (29). Gomez et al. (2022) underscored the resistance of Salmonella isolates in poultry and pork to various antimicrobials, emphasizing the need for control in meat production chains (30).

The studies by Golowczyc et al. in 2007 and 2008

**Table 11.** Calculation of relative expression and statistical analysis of data obtained from Real-time PCR reaction for tetA gene

House (16S rRNA)			
Name	CT	CT	Average ( CT )
Treat	18.74	18.42	18.58
Non-Treat	18.64	18.88	18.76

Gene ( <i>sip</i> )			
Name	CT	CT	Average ( CT )
Sample 1 Treat	19.71	19.85	19.78
Sample 2 Treat	20.33	20.11	20.22
Sample 3 Treat	20.08	19.87	19.975
Sample 1 Non-Treat	18.97	18.91	18.94
Sample 2 Non-Treat	19.03	19.14	19.085
Sample 3 Non-Treat	18.85	19.07	18.96

$\Delta\Delta\text{CT}$ Method			
Name	$\Delta\text{CT}$	$\Delta\Delta\text{CT}$	Fold change
Sample 1 Treat	1.2	1.02	0.493116352
Sample 2 Treat	1.64	1.315	0.401925495
Sample 3 Treat	1.395	1.195	0.436786448
Sample 1 Non-Treat	0.18	-0.055	
Sample 2 Non-Treat	0.325	0.09	
Sample 3 Non-Treat	0.2	-0.035	
<b>Avg Fold change</b>	<b>0.4439428</b>		

T-Test	
P-value	Significant or Not
0.001127783	Significant

emphasized the inhibitory power of Lactobacillus kefir strains against Salmonella, suggesting their potential as probiotics (31, 32). Jr. et al. (2018) demonstrated the probiotic potential of Lactobacillus deliverurans strain Z1 in protecting mice from Salmonella infection (33).

The increase in antibiotic resistance in Salmonella has led to the exploration of alternative approaches to control and prevent enteric bacterial infections. Studies have highlighted the potential role of probiotics, particularly Lactobacillus strains, including Lactobacillus plantarum, Lactobacillus salivarius, Lactobacillus amylovorus, and Lactobacillus kefir, to protect against Salmonella infection through interference with its growth and virulence properties (34, 35). The antagonistic activity of Lactobacillus strains has been shown to inhibit the growth of Salmonella typhimurium in vitro, with significant growth inhibition rates observed. The inhibitory effects are attributed to various factors, including the production of antimicrobial metabolites such as

lactic acid and bacteriocins (36). Lactobacillus strains have been found to exhibit adhesion capacities, auto aggregation, and coaggregation with Salmonella, leading to decreased adherence and invasion of host cells by Salmonella (32). Additionally, the production of antimicrobial metabolites, including lactic acid and bacteriocins, contributes to the inhibition of Salmonella growth and virulence properties. Additionally, research has shown that *Salmonella* spp. isolated from food sources exhibit resistance to various antibiotics, emphasizing the need for alternative control measures (13, 37).

In the context of antibiotic resistance, the *tetA* gene, which confers resistance to tetracycline, and the *sip* gene, associated with Salmonella pathogenicity, have been subjects of interest. The effect of kefir lactococcus in preventing the expression of these genes, as well as its impact on antibiotic resistance in Salmonella, presents a promising avenue for further research and potential application in infection

control and prevention. Furthermore, the diversity of antimicrobial resistance genes in kefir and yogurt, including antibiotic target protection and antibiotic efflux mechanisms, has been investigated, shedding light on the complex interplay between probiotics and antibiotic resistance (35).

In conclusion, *Salmonella typhimurium* poses a considerable risk in causing intestinal infections, with antibiotic resistance and multidrug resistance being major concerns. The use of alternative antibiotics and the potential of probiotic interventions, particularly with lactobacillus strains, are suggested for effective treatment and prevention. The comparison with other studies emphasizes the widespread prevalence of *Salmonella* strains and the urgent need for control measures in various sources, including food, animals, and the environment. The interplay between antibiotic resistance in *Salmonella*, the role of specific genes such as *tetA* and *sip*, and the potential of kefir lactococillus in preventing antibiotic resistance and pathogenicity gene expression represents a dynamic and evolving area of study. Further research in this field holds promise for the development of innovative strategies to address antibiotic resistance and enhance infection control and prevention measures.

#### *Ethical considerations*

In this research, sampling was not done directly from the patient, therefore, there is no interference in the process of diagnosis and treatment of the patient.

#### *Acknowledgments*

The authors would like to thank all individuals whose participation made this study possible.

#### REFERENCES

- Janda JM, Abbott SL. The changing face of the family Enterobacteriaceae (Order: "Enterobacterales"): New members, taxonomic issues, geographic expansion, and new diseases and disease syndromes. *Clinical microbiology reviews* 2021;34(2):10.1128/cmr.00174-20.
- Park J-w, Lee H, Park SY, Kim TH. Epidemiological, clinical, and microbiological characteristics of carbapenemase-producing Enterobacteriaceae bloodstream infection in the Republic of Korea. *Antimicrobial Resistance & Infection Control* 2019;8(1):1-9.
- Jajere SM. A review of *Salmonella enterica* with particular focus on the pathogenicity and virulence factors, host specificity and antimicrobial resistance including multidrug resistance. *Vet World* 2019;12(4):504-21
- Golberg D, Kroupitski Y, Belausov E, Pinto R, Sela S. *Salmonella Typhimurium* internalization is variable in leafy vegetables and fresh herbs. *International journal of food microbiology* 2011;145(1):250-57.
- Cox NA, Cason JA, Richardson LJ. Minimization of *Salmonella* contamination on raw poultry. *Annu Rev Food Sci Technol* 2011;2:75-95.
- Finstad S, O'Bryan CA, Marcy JA, Crandall PG, Ricke

- SC. *Salmonella* and broiler processing in the United States: Relationship to foodborne salmonellosis. *Food Research International* 2012;45(2):789-94.
- Pui C, Wong W, Chai L, et al. *Salmonella*: A foodborne pathogen. *International Food Research Journal* 2011;18(2).
- Feasey NA, Dougan G, Kingsley RA, Heyderman RS, Gordon MA. Invasive non-typhoidal salmonella disease: an emerging and neglected tropical disease in Africa. *Lancet* 2012;379(9835):2489-99.
- Organization WH. *Antimicrobial resistance: global report on surveillance*: World Health Organization, 2014.
- Levy H, Diallo S, Tennant SM, et al. PCR method to identify *Salmonella enterica* serovars Typhi, Paratyphi A, and Paratyphi B among *Salmonella* Isolates from the blood of patients with clinical enteric fever. *J Clin Microbiol* 2008;46(5):1861-6.
- Kumar A, Arora V, Bashamboo A, Ali S. Detection of *Salmonella typhi* by polymerase chain reaction: implications in diagnosis of typhoid fever. *Infect Genet Evol* 2002;2(2):107-10.
- Yang S-M, Kim E, Kim D, et al. Rapid Real-Time Polymerase Chain Reaction for *Salmonella* Serotyping Based on Novel Unique Gene Markers by Pangenome Analysis. *Frontiers in Microbiology* 2021;12.
- D VTN, Venkitanarayanan K, Kollanoor Johny A. Antibiotic-Resistant *Salmonella* in the Food Supply and the Potential Role of Antibiotic Alternatives for Control. *Foods* 2018;7(10).
- Ahn S-I, Kim MS, Park DG, Han BK, Kim YJ. Effects of probiotics administration on lactose intolerance in adulthood: A meta-analysis. *Journal of Dairy Science* 2023;106(7):4489-501.
- Hamilton-Miller J. Probiotics and prebiotics in the elderly. *Postgraduate Medical Journal* 2004;80(946):447-51.
- Tegegne BA, Kebede B. Probiotics, their prophylactic and therapeutic applications in human health development: A review of the literature. *Heliyon* 2022.
- Toh ZQ, Anzela A, Tang ML, Licciardi PV. Probiotic therapy as a novel approach for allergic disease. *Front Pharmacol* 2012;3:171.
- Kim DH, Jeong D, Kim H, Kang IB, Chon JW, Song KY, Seo KH. Antimicrobial Activity of Kefir against Various Food Pathogens and Spoilage Bacteria. *Korean J Food Sci Anim Resour* 2016;36(6):787-90.
- de Oliveira Leite AM, Miguel MA, Peixoto RS, Rosado AS, Silva JT, Paschoalin VM. Microbiological, technological and therapeutic properties of kefir: a natural probiotic beverage. *Braz J Microbiol* 2013;44(2):341-9.
- Colautti A, Orecchia E, Comi G, Iacumin L. Lactobacilli, a Weapon to Counteract Pathogens through the Inhibition of Their Virulence Factors. *J Bacteriol* 2022;204.
- Cappuccino JG, Welsh C. *Microbiology: A Laboratory Manual*: Pearson, 2019.
- Patel JB. *Performance Standards for Antimicrobial Susceptibility Testing: Clinical and Laboratory Standards Institute*, 2017.
- Brenner F, Villar R, Angulo F, Tauxe R, Swaminathan B. *Salmonella nomenclature*. *Journal of clinical microbiology* 2000;38(7):2465-67.
- Clinical, Institute LS. *Performance standards for antimicrobial susceptibility testing: Clinical and Laboratory Standards Institute Wayne, PA, 2017:106-12.*

25. Talib N, Mohamad NE, Yeap SK, et al. Isolation and characterization of *Lactobacillus* spp. from kefir samples in Malaysia. *Molecules* 2019;24(14):2606.
26. Nosrati S, Azar S, Dezfoulian M, Tabarace B, Fallah F. Prevalence of *Salmonella typhimurium*, enteritidis typhimurium serotypes in the foods in Mofid children's Hospital. *Res in Med* 2012;36:43-48.
27. Nazari Moghadam M, Rahimi E, Shakerian A, Momtaz H. Prevalence of *Salmonella Typhimurium* and *Salmonella Enteritidis* isolated from poultry meat: virulence and antimicrobial-resistant genes. *BMC Microbiol* 2023;23(1):168.
28. Ribeiro VB, Lincopan N, Landgraf M, Franco BD, Destro MT. Characterization of class 1 integrons and antibiotic resistance genes in multidrug-resistant *Salmonella enterica* isolates from foodstuff and related sources. *Brazilian Journal of Microbiology* 2011;42:685-92.
29. Dantas ST, Camargo CH, Tiba-Casas MR, et al. Environmental persistence and virulence of *Salmonella* spp. Isolated from a poultry slaughterhouse. *Food Research International* 2020;129:108835.
30. Ruvalcaba-Gómez JM, Villagrán Z, Valdez-Alarcón JJ, et al. Non-antibiotics strategies to control *Salmonella* infection in poultry. *Animals* 2022;12(1):102.
31. Golowczyc MA, Gugliada MJ, Hollmann A, et al. Characterization of homofermentative lactobacilli isolated from kefir grains: potential use as probiotic. *J Dairy Res* 2008;75(2):211-7.
32. Golowczyc MA, Mobili P, Garrote GL, Abraham AG, De Antoni GL. Protective action of *Lactobacillus* kefir carrying S-layer protein against *Salmonella enterica* serovar Enteritidis. *Int J Food Microbiol* 2007;118(3):264-73.
33. Abatemarco Júnior M, Sandes SHC, Ricci MF, Arantes RME, Nunes Á C, Nicoli JR, Neumann E. Protective Effect of *Lactobacillus diolivorans* 1Z, Isolated From Brazilian Kefir, Against *Salmonella enterica* Serovar Typhimurium in Experimental Murine Models. *Front Microbiol* 2018;9:2856.
34. Lya B, Laurence C, Isabelle L-L, Jean-Philippe C. &Lactobacillus& spp. decrease the susceptibility of &Salmonella& Typhimurium to the last resort antibiotic azithromycin. *bioRxiv* 2023:2023.10.01.560186.
35. Gut AM, Vasiljevic T, Yeager T, Donkor ON. Anti-salmonella properties of kefir yeast isolates: An in vitro screening for potential infection control. *Saudi J Biol Sci* 2022;29(1):550-63.
36. Mulaw G, Muleta D, Tesfaye A, Sisay T. Protective Effect of Potential Probiotic Strains from Fermented Ethiopian Food against *Salmonella Typhimurium* DT104 in Mice. *Int J Microbiol* 2020;2020:7523629.
37. Wang X, Biswas S, Paudyal N, Pan H, Li X, Fang W, Yue M. Antibiotic Resistance in *Salmonella Typhimurium* Isolates Recovered From the Food Chain Through National Antimicrobial Resistance Monitoring System Between 1996 and 2016. *Front Microbiol* 2019;10:985.